



Impact of reactive bromine chemistry in the troposphere

R. von Glasow, R. von Kuhlmann, M. G. Lawrence, U. Platt, P. J. Crutzen

► To cite this version:

R. von Glasow, R. von Kuhlmann, M. G. Lawrence, U. Platt, P. J. Crutzen. Impact of reactive bromine chemistry in the troposphere. *Atmospheric Chemistry and Physics*, 2004, 4 (11/12), pp.2481-2497. hal-00295566

HAL Id: hal-00295566

<https://hal.science/hal-00295566>

Submitted on 8 Dec 2004

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Impact of reactive bromine chemistry in the troposphere

R. von Glasow^{1,2}, R. von Kuhlmann^{3,4}, M. G. Lawrence³, U. Platt², and P. J. Crutzen^{2,3}

¹Institute for Environmental Physics, Universität Heidelberg, Germany

²Center for Atmospheric Sciences, Scripps Institution of Oceanography, University of California, San Diego, USA

³Atmospheric Chemistry Division, Max-Planck-Institut für Chemie, Mainz, Germany

⁴now at: German Aerospace Center (DLR), Bonn-Oberkassel, Germany

Received: 28 June 2004 – Published in Atmos. Chem. Phys. Discuss.: 24 August 2004

Revised: 2 December 2004 – Accepted: 7 December 2004 – Published: 8 December 2004



Abstract. Recently several field campaigns and satellite observations have found strong indications for the presence of bromine oxide (BrO) in the free troposphere. Using a global atmospheric chemistry transport model we show that BrO mixing ratios of a few tenths to 2 pmol mol⁻¹ lead to a reduction in the zonal mean O₃ mixing ratio of up to 18% in widespread areas and regionally up to 40% compared to a model run without bromine chemistry. A lower limit approach for the marine boundary layer, that does not explicitly include the release of halogens from sea salt aerosol, shows that for dimethyl sulfide (DMS) the effect is even larger, with up to 60% reduction of its tropospheric column. This is accompanied by dramatic changes in DMS oxidation pathways, reducing its cooling effect on climate. In addition there are changes in the HO₂ : OH ratio that also affect NO_x and PAN. These results imply that potentially significant strong sinks for O₃ and DMS have so far been ignored in many studies of the chemistry of the troposphere.

1 Introduction


In the last two decades reactive halogens have been identified as important reactants in the troposphere (see e.g. von Glasow and Crutzen (2003) for an overview). BrO has been found in the boundary layer during polar ozone depletion events (ODEs) (Hausmann and Platt, 1994), over salt lakes (Hebestreit et al., 1999; Stutz et al., 2002; Hönninger et al., 2004), and in the marine boundary layer (MBL) (Leser et al., 2003; Saiz-Lopez et al., 2004) and also in volcanic plumes (Bobrowski et al. (2003), N. Bobrowski, pers. comm.). Satellite observations showed the widespread presence of BrO in the troposphere outside the polar regions (Wagner and Platt, 1998; Pundt et al., 2000; Fitzenberger et al., 2000;

Wagner et al., 2001; Van Roozendaal et al., 2002; Richter et al., 2002; Hollwedel et al., 2003) with global background vertical columns of about 1–3 × 10¹³ molec cm⁻², corresponding to BrO mixing ratios of 0.5–2 pmol mol⁻¹ if uniformly mixed in the troposphere. Comparisons with balloon and ground based measurements in mid and high northern latitudes (between 42 and 68° N, Harder et al. (1998); Fitzenberger et al. (2000); Van Roozendaal et al. (2002)) as well as the diurnal variation of ground based BrO column measurements (Van Roozendaal et al., 2002) indicated that the tropospheric BrO was mainly located within the free troposphere (FT). Direct evidence for free tropospheric BrO in the Arctic comes from balloon (Fitzenberger et al., 2000) and high-altitude aircraft observations (McElroy et al., 1999). Such observations have not been made yet at lower latitudes. Although most evidence indicates that the measured BrO is located in the FT, these data do not exclude contributions from the MBL. Ongoing work is estimating tropospheric vertical columns of BrO from satellite data but it is expected to yield reliable data only in the polar regions or other regions of strongly elevated BrO (J. Hollwedel, pers. comm.).




Halogens affect ozone by directly destroying it and by reducing its production. The main destruction cycle is (the numbers in parentheses refer to the numbering of the reactions in Table 2):



Uptake of HOBr is an alternative route for reaction (Br16) which cycles Br back to the gas phase. The simplest catalytic ozone production cycle is:



Correspondence to: R. von Glasow
 (Roland.von.Glasow@iup.uni-heidelberg.de)



This is being “shortcut” by cycle I which converts HO_2 back to OH without oxidizing NO to NO_2 . The net effect on odd oxygen ($=\text{O}_x$) is therefore to destroy one molecule of O_x ($=\text{BrO}$) instead of producing one ($=\text{NO}_2$) where O_x is defined based on Crutzen and Schmailzl (1983) as $\text{O}_x = \text{O}_3 + \text{O} + \text{O}(^1\text{D}) + \text{NO}_2 + 2\text{NO}_3 + 3\text{N}_2\text{O}_5 + \text{HNO}_4 + \text{ClO} + 2\text{Cl}_2\text{O}_2 + 2\text{OCIO} + \text{BrO}$. Another “shortcut” occurs in the NO_x cycle:



As both BrO and NO_2 are members of the O_x family, this cycle oxidizes NO to NO_2 without changing the sum of O_x (see also discussion in e.g. Stutz et al. (1999); Platt and Hönninger (2003)). Reaction (Br 9), however, does not lead to a change of the NO to NO_2 ratio on a global scale as explained in Sect. 3.4.

In this study we examine the potential effects of the presence of $0.5\text{--}2\text{ pmol mol}^{-1}$ of BrO on the photochemistry (mainly ozone and DMS) in the (free) troposphere and discuss where we should focus our future research in this field. We describe the model and the source scenarios that we used in Sect. 2 and will discuss the implications for tropospheric chemistry in Sect. 3 focussing on O_3 , HO_x , NO_x , and DMS. In Sect. 4 we conclude and list future research needs.

2 Model description and source scenarios


For this study we used the three-dimensional chemical transport model MATCH-MPIC (Lawrence et al., 1999; von Kuhlmann et al., 2003) which includes a comprehensive treatment of tropospheric gas phase chemistry including non-methane hydrocarbons. We added reactions relevant for bromine processing and bromine-related ozone destruction and the oceanic sources of DMS after Kettle et al. (1999) and its destruction by OH, NO_3 , BrO, and Br. The additional reactions are listed in Table 2. Wet and dry deposition are included as a function of the Henry’s law constant of the respective compounds.

The actual sources for the tropospheric inorganic bromine (Br_{inorg}) are still poorly constrained. They include downward transport from the stratosphere in mid to high latitudes, “spillout” and uplifting from polar surface ODEs, upward transport from the marine boundary layer, volcanoes, organohalogens (e.g. CH_3Br , CH_2Br_2 , CHBr_3 , and CHBr_2Cl) which are broken down by photolysis and/or reaction with OH, release of inorganic halogens by biomass burning as shown in laboratory experiments (B. Keene, pers. comm.), or other, so far unidentified processes. Many of

these sources will have a seasonal cycle which we do not take into account in this exploratory study. To sustain a tropospheric mixing ratio of Br_{inorg} of about 3 pmol mol^{-1} and assuming a Br_{inorg} lifetime of 1–2 weeks in the FT a global source of $2.5\text{--}5 \times 10^{-18}\text{ mol mol}^{-1}\text{ s}^{-1}$ or about $5.6\text{--}11 \times 10^7\text{ molec cm}^{-2}\text{ s}^{-1}$ would be needed. It is likely that the total source is actually composed of several of the aforementioned sources.

To reproduce the previously mentioned mixing ratios of BrO in the FT we tested four different hypothetical source scenarios (see Table 1): “tropics”, “high lat”, “const”, and “strat” which reflect different possible source types/regions, namely the tropics, high latitudes, a source constant with latitude, and only downward flux of inorganic bromine from the stratosphere plus photochemical breakdown of CH_3Br . We have chosen this set of model runs to study the differences in the impacts related to these spatial distributions of BrO. Except for the case “strat” we do not differentiate in any of these scenarios between the possible sources that we have mentioned above. As no good global data on the different source processes is available we cannot derive source scenarios that explicitly take the spatial coverage of single sources into account. Furthermore, the scarcity of observational data does not allow us to choose one of these scenarios as a “best guess”.

For the scenario “tropics” we used the simplified reaction:



with a generic organic bromine compound Br_{org} (with fixed mixing ratio of 30 pmol mol^{-1} Br) as bromine source, constituting a bromine source of about $100\text{ molec cm}^{-3}\text{ s}^{-1}$ at a temperature of $T=250\text{ K}$ and an OH concentration of $10^6\text{ molec cm}^{-3}$. The OH dependence ensures a direct coupling with the photochemistry and makes this bromine source most important in the tropics where OH concentrations are highest.

The bromine source of the scenario “high lat” is independent of photochemistry and increasing with latitude, corresponding to a dominance of downward transport from the stratosphere and spillout from polar surface ODEs. We used $F_{\text{high lat}} = (15 + 0.5\phi) \frac{[M(\phi)]}{[M_0(\phi)]}\text{ molec cm}^{-3}\text{ s}^{-1}$, where ϕ is the latitude in degrees and $[M]$ and $[M_0]$ the concentration of air molecules at current altitude and at the surface, respectively.

In the scenario “const” the source is also independent of photochemistry but constant with latitude, $F_{\text{const}} = 34 \frac{[M(\phi)]}{[M_0(\phi)]}\text{ molec cm}^{-3}\text{ s}^{-1}$. In scenarios “high lat” and “const” no time dependence of the bromine source was assumed, whereas the dependence on [OH] in scenario “tropics” leads to a diurnally and seasonally varying bromine source.

The scaling with the concentration of air molecules in scenarios “high lat” and “const” ensures that the Br_{inorg} source is constant with altitude in mixing ratio ($\text{mol mol}^{-1}\text{ s}^{-1}$) but obviously not in concentration ($\text{molec cm}^{-3}\text{ s}^{-1}$). Due to different chemical regimes and washout rates the impact and

Table 1. Overview of scenarios used in this study.

scenario	heterogeneous bromine reactions	description
high lat	$k'_{het}=10. \times k_{het}(\gamma=0.1)^a$	Br source increases with latitude
high lat, slow recyc.	$\gamma=0.1$	Br source increases with latitude
high lat, no recyc.	off	Br source increases with latitude
const	$k'_{het}=10. \times k_{het}(\gamma=0.1)^a$	Br source constant with latitude
tropics	$k'_{het}=10. \times k_{het}(\gamma=0.1)^a$	Br source decreases with latitude
strat	$k'_{het}=10. \times k_{het}(\gamma=0.1)^a$	Br downward transport from stratosphere plus decomposition of CH ₃ Br
nohal	off	no bromine

^a see explanation in text, the effective gamma for these runs is approx. $\gamma=0.5$.

Table 2. Bromine reactions.



no	reaction	<i>n</i>	<i>A</i> [(cm ⁻³) ^{1-<i>n</i>} s ⁻¹]	$-E_a/R$ [K]	Reference
Br 1	$\text{Br} + \text{O}_3 \rightarrow \text{BrO} + \text{O}_2$	2	1.7×10^{-11}	-800	Sander et al. (2003)
Br 2	$\text{Br} + \text{HO}_2 \rightarrow \text{HBr} + \text{O}_2$	2	1.5×10^{-11}	-600	Sander et al. (2003)
Br 3	$\text{Br} + \text{C}_2\text{H}_4 \xrightarrow{\text{O}_2} \text{HBr} + \text{C}_2\text{H}_5\text{O}_2$	2	$5. \times 10^{-14}$		Singh and Zimmerman (1992)
Br 4	$\text{Br} + \text{HCHO} \xrightarrow{\text{O}_2} \text{HBr} + \text{CO} + \text{HO}_2$	2	1.7×10^{-11}	-800	Sander et al. (2003)
Br 5	$\text{Br} + \text{CH}_3\text{OOH} \rightarrow \text{CH}_3\text{OO} + \text{HBr}$	2	2.66×10^{-12}	-1610	Mallard et al. (1993)
Br 6	$\text{Br} + \text{BrNO}_3 \rightarrow \text{Br}_2 + \text{NO}_3$	2	4.9×10^{-11}		Orlando and Tyndall (1996)
Br 7	$\text{BrO} + \text{HO}_2 \rightarrow \text{HOBr} + \text{O}_2$	2	3.4×10^{-12}	540	Sander et al. (2003)
Br 8	$\text{BrO} + \text{CH}_3\text{OO} \rightarrow 0.72 \text{ HOBr} + 0.28 \text{ Br} + 0.28 \text{ HO}_2 + \text{HCHO}^a$	2	5.7×10^{-12}		Aranda et al. (1997)
Br 9	$\text{BrO} + \text{NO} \rightarrow \text{Br} + \text{NO}_2$	2	8.8×10^{-12}	260	Sander et al. (2003)
Br 10	$\text{BrO} + \text{NO}_2 \xrightarrow{M} \text{BrNO}_3$	2	^b		Sander et al. (2003)
Br 11	$\text{BrO} + \text{BrO} \rightarrow 1.62 \text{ Br} + 0.19 \text{ Br}_2 + \text{O}_2$	2	2.95×10^{-12}	40	based on Sander et al. (2003), 2 channels (different T-dependence) combined
Br 12	$\text{HBr} + \text{OH} \rightarrow \text{Br} + \text{H}_2\text{O}$	2	1.1×10^{-11}		Sander et al. (2003)
Br 13	$\text{BrNO}_3 \rightarrow \text{BrO} + \text{NO}_2$	1	^b		Orlando and Tyndall (1996)
Br 14	$\text{BrO} + h\nu \xrightarrow{\text{O}_2} \text{Br} + \text{O}_3$	1	^c		DeMore et al. (1997)
Br 15	$\text{Br}_2 + h\nu \rightarrow 2 \text{ Br}$	1	^c		Hubinger and Nee (1995)
Br 16	$\text{HOBr} + h\nu \rightarrow \text{Br} + \text{OH}$	1	^c		Ingham et al. (1999)
Br 17	$\text{BrNO}_3 + h\nu \rightarrow \text{Br} + \text{NO}_3$	1	^c		DeMore et al. (1997)
Br 18	$\text{HOBr} + \text{HBr} \xrightarrow{het} \text{Br}_2 (+ \dots)$	2			see text
Br 19	$\text{BrNO}_3 \xrightarrow{het} \text{HOBr} (+ \dots)$	1			see text
Br 20	$\text{Br}_{org} + \text{OH} \rightarrow \text{Br} (+ \dots)$	2	1.7×10^{-11}	-1215	assumed, only for “tropics”, [Br _{org}]=30 pmol mol ⁻¹
Br 21	$\text{CH}_3\text{Br} + \text{OH} \rightarrow \text{Br} (+ \dots)$	2	1.7×10^{-12}	-1215	Atkinson et al. (2003), only for “strat”
S1	$\text{DMS} + \text{OH} \xrightarrow{\text{O}_3} (\text{CH}_3\text{SO}_2 +) \text{HCHO}^d$	2	1.12×10^{-11}	-253	after Atkinson et al. (1997)
S2	$\text{DMS} + \text{OH} \rightarrow (0.5\text{DMSO} + 0.5\text{SO}_2 +) 0.5 \text{HO}_2 + \text{CH}_3\text{OO}^d$	2	^b		after Atkinson et al. (1997)
S3	$\text{DMS} + \text{NO}_3 \xrightarrow{\text{O}_3} (\text{CH}_3\text{SO}_2 +) \text{HNO}_3 + \text{HCHO}^d$	2	1.9×10^{-13}	520	Atkinson et al. (1999)
S4	$\text{DMS} + \text{Br} \xrightarrow{\text{O}_3} (\text{CH}_3\text{SO}_2 +) \text{HBr} + \text{HCHO}^d$	2	9.0×10^{-11}	-2386	Jefferson et al. (1994)
S5	$\text{DMS} + \text{BrO} \rightarrow (\text{DMSO} +) \text{Br}$	2	2.54×10^{-14}	850	Ingham et al. (1999)

n is the order of the reaction. ^a two reactions combined, ^b special rate functions (pressure dependent), ^c photolysis rates calculated online, ^d simplified final products of importance for HO_x chemistry included. The rate coefficients are calculated with $k=A \times \exp(-\frac{E_a}{RT})$. Note that the breakdown products of DMS are ignored in the model.

lifetime and therefore “steady state” mixing ratio is dependent on altitude and latitude.

The only bromine sources in scenario “strat” are downward flux of inorganic bromine from the stratosphere and decomposition of CH_3Br (with a constant mixing ratio of 10 pmol mol^{-1}) in the troposphere by reaction with OH. To calculate the downward flux we used fixed concentrations of inorganic bromine in the stratosphere (dependent on latitude and season) as a boundary condition, the data are taken from a two-dimensional stratospheric model (Brühl et al., 1998; WMO, 2003). As explained by von Kuhlmann et al. (2003) a factor of 0.5 has to be used to improve the stratosphere-troposphere exchange of O_3 in the model. In the scenario “strat” we applied the same ratio for the downward flux of inorganic bromine.

If only gas phase reactions are included, the cycling of HOBr, HBr, and BrNO_3 is rather slow and the resulting BrO concentrations are small. It was shown in many studies, however, that very efficient cycling of inorganic bromine occurs on and within aerosol particles (see e.g. overview by von Glasow and Crutzen (2003) and references therein). Our simplified approach for the heterogeneous reactions is:



based on detailed reaction cycles (Fan and Jacob, 1992; Abbatt, 1994; Vogt et al., 1996; Sander et al., 1999; Fickert et al., 1999). In the current model version no aqueous phase species are considered, therefore we listed only the gas phase products, other products are assumed to be taken up irreversibly by the aerosol. We used the approach of the reaction of HOBr with HBr on aerosol surfaces because of the high solubility of HBr, a possible enrichment of bromide on the aerosol surface (Jungwirth and Tobias, 2002), and the already mentioned high reaction probabilities on aerosols. Note that reaction (Br19) constitutes loss of NO_x for the gas phase (see discussion in Sect. 3.4). We used the heterogeneous reaction rates as calculated by Dentener and Crutzen (1993) (based on Langner and Rodhe (1991)) that were derived based on climatological monthly mean values for all meteorological parameters (e.g. relative humidity), a reaction probability of $\gamma=0.1$, and taking gas phase diffusion limitations into account. To investigate the implications of bromine recycling on aerosol we repeated scenario “high lat” without aerosol recycling (“no recycling”) and a 10 times faster recycling rate ($k'_{\text{het}}=10 \times k_{\text{het}}(\gamma=0.1)$) which approximately corresponds to $\gamma=0.5$ for the size distribution chosen by Dentener and Crutzen (1993). Very high reaction probabilities of the involved bromine species have been observed on different substrates (Sander et al., 2003), and, as explained below, the results with faster heterogeneous reaction rates yield better agreement with the available information on BrO tropospheric vertical columns, therefore we used the higher re-

cycling rate k'_{het} for all scenarios except for “high lat, no recycling” and “high lat, slow recycling”.

Many details of the bromine recycling are still unknown, cycling of HOBr on frozen salt surfaces (Adams et al., 2002) or cirrus clouds, or in stratiform clouds (von Glasow et al., 2002b) are additional routes. If these bromine recycling reactions or other processes that reduce the loss of inorganic bromine are confirmed to be of importance, a smaller source for inorganic bromine would be needed to reproduce the reported BrO mixing ratios.

As already mentioned, the actual recycling of inorganic halogens on aerosols happens in a more complicated way but by using this approach we assume a dependence on the available aerosol surface area and yield BrO mixing ratios that are close to those deduced from satellite observations and balloon measurements (see Sect. 3.1). No aerosol components are transported in the current model version, so potential accumulation effects on the aerosol or a spatial redistribution of bromine by transport/settling of aerosol and time-delayed release of inorganic bromine from the aerosol cannot be simulated in this model version.

A model resolution of $11.25^\circ \times 11.25^\circ$ at the equator (T10) with 28 vertical levels is used. This is sufficiently high to examine the large scale features and the overall effects of bromine especially considering the uncertainties in the source strength and distribution. All scenarios were run for 12 months after a spinup of 16 months each.

3 Results

3.1 Br_{inorg} distributions

We start by briefly discussing the distribution of bromine species in the different scenarios before we explain the impacts of bromine on other trace gases focusing on the free troposphere. As explained in the previous section we use artificial in situ sources for bromine in order to analyze the atmospheric response based on different latitudinal distributions of bromine. These sources do not explicitly take sea salt aerosol into account, therefore the bromine levels in the boundary layer in all runs – and all related consequences for photochemistry – are lower limits. This setup was chosen because the main purpose of this study is to investigate effects in the free troposphere and to provide lower limit estimates for the boundary layer. Ongoing work deals with the explicit treatment of sea salt aerosol chemistry on a global scale.

Total tropospheric Br_{inorg} levels in all scenarios are about $1\text{--}6 \text{ pmol mol}^{-1}$ (see e.g. Fig. 1 for scenario “high lat”) except for run “strat” where they are below 1 pmol mol^{-1} . In scenarios with a faster recycling rate compared to runs with slower or no recycling Br_{inorg} increases by more than 200% in regions with high wet deposition rates, i.e. the tropics but also in southern mid latitudes in altitudes of 900 to 500 hPa. Faster recycling implies higher BrO : Br_{inorg} and




Fig. 1. Zonally and annually averaged mean of the Br_{inorg} mixing ratio (in pmol mol^{-1}) resulting from our source scenario “high lat”. The ordinate is the σ -level multiplied by 1000 which is approximately the pressure in hPa and the abscissa is latitude in degrees. Note that 24h averages are plotted. Seasalt aerosol is not explicitly included as a source therefore boundary layer values are lower limits.




Fig. 2. Zonally and annually averaged mean of the ratio of BrO to Br_{inorg} resulting from our source scenario “high lat”. The ordinate is the σ -level multiplied by 1000 which is approximately the pressure in hPa and the abscissa is latitude in degrees. Note that 24h averages are plotted.

$\text{Br}_2 : \text{Br}_{\text{inorg}}$ values and therefore less deposition as these two species are relatively insoluble. HBr and BrNO_3 (and HOBr), on the other hand, get washed out rapidly. In the “tropics” scenario Br_{inorg} mixing ratios in the tropics are

up to 5 pmol mol^{-1} and only around 2 pmol mol^{-1} in high latitudes whereas in the case “high lat” they are less than 2 pmol mol^{-1} in the tropics and up to 5 pmol mol^{-1} in high latitudes. In the scenario “const” the numbers are similar




Fig. 3. Zonally and annually averaged mean of the lifetime of Br_{inorg} (in days) resulting from our source scenario “high lat”. The ordinate is the σ -level multiplied by 1000 which is approximately the pressure in hPa and the abscissa is latitude in degrees.




Fig. 4. Zonally and annually averaged mean of the BrO mixing ratio (in pmol mol^{-1}) resulting from our source scenario “high lat”. The ordinate is the σ -level multiplied by 1000 which is approximately the pressure in hPa and the abscissa is latitude in degrees. Note that 24h averages are plotted. Seasalt aerosol is not explicitly included as a source therefore boundary layer values are lower limits.

to “high lat” with a smaller but discernable increase from the tropics to higher latitudes (see explanation below). In all scenarios Br_{inorg} mixing ratios increase with altitude (see below). Assuming photochemical steady state we calculated the lifetime of Br_{inorg} that is between 10 and 25 days for the

free troposphere in case “high lat, fast recycling” (see Fig. 3). The influence of the recycling of bromine on aerosols on the Br_{inorg} lifetime is decreasing with altitude and is strongest below about 400 hPa. In case “fast recycling” the Br_{inorg} lifetime is about 20% to 130% higher than in case “slow




Fig. 5. Zonally and annually averaged mean of the BrO mixing ratio (in pmol mol^{-1}) resulting from our source scenario “tropics”. The ordinate is the σ -level multiplied by 1000 which is approximately the pressure in hPa and the abscissa is latitude in degrees. Note that 24h averages are plotted. Seasalt aerosol is not explicitly included as a source therefore boundary layer values are lower limits.




Fig. 6. Zonally and annually averaged mean of the BrO mixing ratio (in pmol mol^{-1}) resulting from our source scenario “strat”. The ordinate is the σ -level multiplied by 1000 which is approximately the pressure in hPa and the abscissa is latitude in degrees. Note that 24h averages are plotted.

recycling” and even 50% to 230% higher than in case “no recycling”.

The zonally averaged annual mean mixing ratio of BrO (24 h average) is less than $0.5 \text{ pmol mol}^{-1}$ in all scenarios

for most of the troposphere and reaches 2 pmol mol^{-1} only in the upper troposphere especially in scenario “tropics” (see Figs. 4, 5, and 6). It increases with altitude and southern latitude. The BrO distribution in the “const” and “tropics”




Fig. 7. Annually averaged tropospheric vertical column of BrO (in molec cm⁻²) for the scenario “high lat”. We used the WMO definition for the determination of the tropopause. Note that 24h averages are plotted. The ordinate is latitude in degrees and the abscissa is longitude in degrees.

runs is latitudinally more uniform than that of case “high lat”. This latitudinal pattern in the scenario “high lat” follows roughly the trend in Br_{inorg} but the zonally averaged ratio of BrO to Br_{inorg} (24 h average) also increases with altitude and with latitude (in all runs) with values of 0.05 near the surface increasing to 0.3 at the top of the troposphere (see Fig. 2). This is mainly due to an increase in solar radiation and therefore shorter lifetimes of HOBr and BrNO_3 . This trend is present in all scenarios but the increase with altitude is faster in runs with higher recycling rates because the faster recycling increases the lifetime of Br_{inorg} . Another factor is the increase in $\text{Br}_2:\text{Br}_{\text{inorg}}$ and a decrease in $\text{HOBr}:\text{Br}_{\text{inorg}}$ with altitude which increases the relative contribution of BrO because the photolysis rate for Br_2 is greater than that for HOBr and increases Br_{inorg} with altitude by a reduction in the washout of Br_{inorg} because of differences in solubility of Br_2 and HOBr.

It has been shown with a one-dimensional model (von Glasow et al., 2002a) that the difference of the wavelength dependence of $\text{O}_3 \rightarrow \text{O}(^1\text{D})$ and Br_2 photolysis leads to a diurnal variation of BrO with small morning and afternoon peaks if HO_2 is the main sink, and a broad diurnal variation with a small peak at noon if NO_2 is the main sink. These two types of diurnal variations are found in our 3D model results as well.

Figures 7 and 8 show the vertically integrated tropospheric column density (using the WMO definition of the tropopause) of BrO for the scenarios “high lat” and “tropics”, respectively, with values of about $0.5\text{--}1 \times 10^{13}$ molec cm⁻² in the tropics and up to 2.4×10^{13} molec cm⁻² in higher latitudes. The large gradient in the tropopause height in the subtropics leads to a maximum in the vertical column of BrO in that region. The maximum values are reached only in the subtropics, whereas in most parts of the troposphere the model BrO vertical columns are somewhat smaller than the tropospheric vertical BrO columns of $<1\text{--}3 \times 10^{13}$ molec cm⁻² that were derived from comparisons of satellite, balloon, and ground measurements (Pundt et al., 2000; Van Roozendael et al., 2002; Richter et al., 2002). For this reason, the results presented here should be regarded as a conservative estimate of the overall effects of bromine chemistry in the free troposphere and, as mentioned before, a lower limit for the boundary layer.

The inorganic bromine levels in scenario “strat” are by far too small to explain the observed tropospheric vertical columns which in this run are only $0.5\text{--}2 \times 10^{12}$ molec cm⁻². The annually and zonally averaged 24 h mixing ratios of BrO are less than $0.25 \text{ pmol mol}^{-1}$ in the troposphere (see Fig. 6). This implies that downward flux of stratospheric inorganic bromine and decomposition of CH_3Br are not sufficient as




Fig. 8. Annually averaged tropospheric vertical column of BrO (in molec cm⁻²) for the scenario “tropics”. We used the WMO definition for the determination of the tropopause. Note that 24h averages are plotted. The ordinate is latitude in degrees and the abscissa is longitude in degrees.

source for free tropospheric inorganic bromine. They would either have to be increased by about a factor of 5–7 or additional sources of reactive bromine have to be active.

3.2 Effect on ozone

The runs with bromine levels that reproduce the observations best lead to a notable decrease in O₃. Even in the runs with smallest BrO mixing ratios of only a few tenths of a pmol mol⁻¹ (scenario “high lat, no recycling” and “strat”, see Fig. 11) the annually and zonally averaged difference to the run without bromine chemistry is of the order of several percent. In the case “high lat, slow recycling” with higher BrO mixing ratios it is already 5–10% throughout the troposphere, whereas in the run with BrO mixing ratios that are closest to the values deduced from satellite and balloon observations (“high lat”) the difference in zonal mean O₃ mixing ratio compared to the run without bromine chemistry is 6–18% (see Fig. 9). Maximum regional differences are up to 40% in the austral summer high latitudes. These numbers imply that there might be an important O₃ destruction mechanism in the troposphere that has been neglected so far.

Even though the mixing ratio and distribution of Br_{inorg} and BrO differ among the different scenarios, the overall vertical and latitudinal distribution of the effect on O₃ remains

Table 3. Tropospheric burdens of O₃ and DMS in Tg.

scenario	O ₃ burden	DMS burden
high lat	157.0	4.612
tropics	147.5	4.787
strat	172.2	5.848
nohal	174.6	6.249

similar in all discussed runs, namely that O₃ destruction is strongest in the FT of the southern hemisphere. It is smallest near the ground in northern mid latitudes. This is due to smaller sensitivities of the photochemistry to perturbations when high O₃ sources and sinks are present as is the case in the polluted regions compared to a greater sensitivity in the more pristine FT in the southern hemisphere where photochemical sources and sinks for O₃ are smaller. In scenario “tropics” the maximum of the difference in zonally averaged O₃ (up to 20% less O₃ compared to a run without bromine chemistry, see Fig. 10) is shifted from higher latitudes towards 50–30° S and strongest in magnitude compared to all other presented cases. Nevertheless, even in this case the main pattern with highest differences in O₃ in the southern FT remains the same.




Fig. 9. Ratio of O_3 in scenario “high lat” to O_3 in scenario “no hal”. The numbers are zonally and annually averaged. The ordinate is the σ -level multiplied by 1000 which is approximately the pressure in hPa and the abscissa is latitude in degrees. Note that 24h averages are plotted.




Fig. 10. Ratio of O_3 in scenario “tropics” to O_3 in scenario “no hal”. The numbers are zonally and annually averaged. The ordinate is the σ -level multiplied by 1000 which is approximately the pressure in hPa and the abscissa is latitude in degrees. Note that 24h averages are plotted.

Compared to the run without bromine chemistry, the tropospheric burden of O_3 (see Table 3) is reduced in scenarios “high lat”, “tropics”, and “strat” by 10%, 15%, and 3%, respectively. The ozone loss by reaction (Br7) equals approximately 28% of the dry deposition in run “high lat”.

3.3 Effect on HO_x

The reaction $\text{BrO} + \text{HO}_2 \longrightarrow \text{HOBr}$ destroys HO_2 and the subsequent photolysis of HOBr releases OH . Both processes are acting in the direction of an increase of the OH to HO_2




Fig. 11. Ratio of O_3 in scenario “strat” to O_3 in scenario “no hal”. The numbers are zonally and annually averaged. The ordinate is the σ -level multiplied by 1000 which is approximately the pressure in hPa and the abscissa is latitude in degrees. Note that 24 h averages are plotted.





Fig. 12. Ratio of $\text{OH}:\text{HO}_2$ in scenario “high lat” to $\text{OH}:\text{HO}_2$ in scenario “no hal”. The numbers are zonally and annually averaged. The ordinate is the σ -level multiplied by 1000 which is approximately the pressure in hPa and the abscissa is latitude in degrees. Note that 24h averages are plotted.

ratio. This shift is highest in the upper FT with up to 50% and 40% in the southern and northern high latitudes, respectively, in the scenario “high lat” (compared to the ratio in the run without bromine, see Fig. 12). In mid and low latitudes this effect is less than 10%. As mentioned in the introduction,


this shift implies less O_3 production because in the reactions



HO₂ gets cycled back to OH without oxidizing NO to NO₂. Furthermore reaction Br7 destroys O_x because BrO is part of the odd oxygen family. The decrease in HO₂ concentrations compared to scenario “no hal” is up to 10% in runs “high lat” and “tropics” with the strongest effect in the free troposphere. In the run “high lat” it is most pronounced at high latitudes whereas it is more uniform with latitude in the run “tropics”.

The OH concentration increases in the upper FT by more than 20% compared to the run without bromine chemistry but this is contrasted by small relative decreases in OH in the tropics. Most OH, however, is located in the tropics where the reduction in O₃ is strong enough to reduce OH, so that the change in global mean OH concentrations (calculated weighted with CH₄ as well as with air mass after Lawrence et al. (2001)) is only for the “tropics” scenario on the order of 1–2%. For the other runs it is virtually indiscernible from scenario “no hal”. The lifetime of CH₄ is almost unchanged compared to “no hal” in all runs but “tropics” where it increased from 9.62 years to 9.83 years, again only 2% so that these effects are negligible.

H₂O₂ decreases with latitude and altitude compared to the run without bromine chemistry (see Fig. 13). This is caused by the decrease in HO₂ which is the main precursor for H₂O₂. The decrease in H₂O₂ in the tropics is a few percent in scenario “high lat” and 5–10% in scenario “tropics” but more than 20% in the high latitude free troposphere. The reduction is therefore larger than that in HO₂, because the formation rate of H₂O₂ in reaction



is quadratic in HO₂. The decrease in H₂O₂ can be of importance for atmospheric chemistry because the gas phase is the main source for aqueous H₂O₂ which is the most important oxidant for S(IV) to S(VI) in the aqueous phase. A reduction in H₂O₂ would reduce the formation rates of particulate sulfate which, however, can possibly be compensated by HOBr (and HOCl) as aqueous phase oxidants as explained e.g. in von Glasow and Crutzen (2004).

3.4 Effect on nitrogen oxides

The runs including bromine chemistry also show changes in the NO_x mixing ratios in mid and high latitudes and especially in the FT. In the scenario “high lat” NO_x concentrations are up to 25% smaller than in the run without bromine chemistry. In the lower troposphere, especially in the tropics, NO_x increases by a few percent. These changes are approximately correlated with changes in HO₂ and anticorrelated with changes in peroxy acetyl nitrate (PAN) and OH. This is caused by an increase in the formation of PAN from the peroxy acetyl (PA) radical which is caused by a decrease in HO₂ that destroys PA.

A decrease in NO_x concentrations reduces the photochemical production of O₃. This highlights that the difference in

O₃ in our model runs is a combination of increased O₃ destruction by bromine and reduced O₃ production due to less NO_x (indirectly caused by bromine) stressing the importance to consider all elements of the chemical system.

As already mentioned in the introduction, the reaction BrO + NO → Br + NO₂ reduces the production rate of O₃ by oxidizing NO to NO₂ without changing the sum of odd oxygen. In the model results, however, the ratio of NO to NO₂ decreases only in the uppermost troposphere in high latitudes and in the stratosphere. In most of the troposphere the ratio increases compared to a run without bromine chemistry. (see Fig. 14). This is caused by decreases in the O₃ and HO₂ concentrations which are crucial for the steady state ratio of NO to NO₂ and which cannot be compensated by the presence of less than 2 pmol mol^{−1} BrO (see e.g. Brasseur et al. (1999)):

$$\frac{[\text{NO}]}{[\text{NO}_2]} = j(\text{NO}_2) \times \left(k_{\text{NO}+\text{O}_3}[\text{O}_3] + k_{\text{HO}_2+\text{NO}}[\text{HO}_2] + k_{\text{CH}_3\text{O}_2+\text{NO}}[\text{CH}_3\text{O}_2] + k_{\text{BrO}+\text{NO}}[\text{BrO}] \right)^{-1} \quad (1)$$

The ratio of BrNO₃ to Br_x is highest in northern mid latitudes near the ground, the highest BrNO₃ mixing ratios are found near the tropopause, they are less than 0.5 pmol mol^{−1} in most of troposphere in all runs except for “tropics” where they can reach up to 1 pmol mol^{−1} in the tropical mid troposphere. The contribution of BrNO₃ to NO_y is less than 0.5%. The heterogeneous reaction of BrNO₃ produces particulate NO₃[−] (see Sander et al. (1999)) and therefore contributes to the loss of NO_x. This NO_x loss by the heterogeneous reaction of BrNO₃ is about 4.5 and 10% of the NO_x loss by reaction NO₂ + OH → HNO₃ in the runs “high lat” and “tropics”, respectively.

3.5 Effect on DMS

Another very important result of our study is the impact of BrO on DMS chemistry with a strong reduction of its mixing ratio and a drastic change in its oxidation pathways. DMS is emitted by marine organisms and is the main natural source for reactive sulfur in the marine atmosphere. The main sinks of DMS are usually thought to be reaction with OH and NO₃, however, the reaction of DMS with BrO can be a significant sink as well (Toumi, 1994). In the cloudy MBL an increase in DMS oxidation by BrO leads to the production of more DMSO which is rapidly taken up by clouds. This reduces the DMS to SO₂ conversion efficiency, which is indicative of the potential for formation of new cloud condensation nuclei (CCN), but increases the size of already present CCN by the formation of particulate sulfur (methyl sulfonic acid and sulfate) (von Glasow et al., 2002b; von Glasow and Crutzen, 2004). This and other feedbacks like increased drizzle formation due to larger CCN would lead to a reduction in cloud




Fig. 13. Ratio of H_2O_2 in scenario “high lat” to H_2O_2 in scenario “no hal”. The numbers are zonally and annually averaged. The ordinate is the σ -level multiplied by 1000 which is approximately the pressure in hPa and the abscissa is latitude in degrees. Note that 24 h averages are plotted.




Fig. 14. Ratio of $\text{NO}:\text{NO}_2$ in scenario “high lat” to $\text{NO}:\text{NO}_2$ in scenario “no hal”. The numbers are zonally and annually averaged. The ordinate is the σ -level multiplied by 1000 which is approximately the pressure in hPa and the abscissa is latitude in degrees. Note that 24 h averages are plotted.

albedo. Boucher et al. (2003) performed a sensitivity study with a global model assuming a constant daytime BrO mixing ratio of 1 pmol mol^{-1} in the MBL and found a reduction of the tropospheric DMS burden by about 30%.

In our model runs the tropospheric DMS burden is reduced due to the reaction $\text{DMS} + \text{BrO}$ by 26%, 23%, and 6% in scenarios “high lat”, “tropics”, respectively (see Table 3). The vertical columns of DMS mixing ratios (dominated by the MBL) are reduced by 3–30% in run “high lat,




Fig. 15. Ratio of the vertical column of DMS for the scenario “high lat” to scenario “no hal” (annually averaged). The ordinate is latitude in degrees and the abscissa is longitude in degrees. Seasalt aerosol is not explicitly included as a source for BrO therefore the reductions in DMS are lower limits.




Fig. 16. Ratio of the vertical column of DMS for the scenario “tropics” to scenario “no hal” (annually averaged). The ordinate is latitude in degrees and the abscissa is longitude in degrees. Seasalt aerosol is not explicitly included as a source for BrO therefore the reductions in DMS are lower limits.

no recycling” with the smaller values close to the equator. In runs “high lat” and “tropics” the difference to “no hal” in many regions is more than 10%, and it is especially high in the regions with highest DMS fluxes and mixing ratios, the Southern Ocean (Kettle et al., 1999) as evident from

Figs. 15 and 16. The spatial distribution of the DMS differences in the model is caused by a latitudinal gradient in BrO with annually averaged 24h mean mixing ratios in the boundary layer of 0.05 (0.4) pmol mol^{-1} in the tropics and 0.3 (0.1) pmol mol^{-1} in high latitudes for scenario “high lat

(“tropics”). Note that DMS mixing ratios are highest in the MBL where our bromine source approach will likely underestimate BrO mixing ratios because we did not include explicitly sea salt aerosol as a source for reactive halogens, so these results are lower limits. The main conclusion from this result is that the global importance of bromine chemistry for the oxidation of DMS should be investigated in more detail.

The main difference of our study to that by Boucher et al. (2003) is, that they assumed a constant daytime BrO mixing ratio of 1 pmol mol^{-1} everywhere in the MBL whereas in our model runs the 24 h average mixing ratio of BrO in the MBL is only $0.1\text{--}0.3 \text{ pmol mol}^{-1}$ in “high lat” and less than $0.05 \text{ pmol mol}^{-1}$ in runs “high lat, no recycling” and “strat”. Again we want to stress that these BrO mixing ratios have to be regarded as lower limits.

The previously discussed increases in DMS oxidation, the shift in its oxidation products, and the changes in resulting new particle formation and growth of existing aerosol particles have the potential to dramatically alter our understanding of the DMS – CCN – climate connection. Moreover, some estimates of the DMS flux from the ocean rely on a comparison with atmospheric models to calculate the photochemical loss rate of DMS without taking oxidation by BrO into account, implying that DMS fluxes from these studies might be underestimating the real fluxes (see also discussion in von Glasow and Crutzen (2004)).

4 Conclusions and future research needs

Our results show that even small levels of reactive bromine can greatly disturb tropospheric chemistry. Bromine related reactions strongly reduce ozone mixing ratios by up to 18% in zonal, annual mean values and up to 40% regionally. Furthermore, they also change the mixing ratios and speciation of the HO_x and NO_x chemical families. This highlights that both the photochemical production and destruction of O_3 is perturbed by the presence of reactive bromine. A lower limit approach for BrO in the marine boundary layer, that does not explicitly treat the release of halogens from sea salt aerosol, showed that the mixing ratios of DMS in the MBL are strongly reduced due to the presence of BrO which in turn also leads to changes in the final products of the sulfur oxidation with potentially drastic changes for the cloud condensation nuclei population in the MBL.

While there is a substantial body of evidence for the presence of relevant levels of tropospheric BrO this needs to be supported with more direct and indirect measurements in all parts of the troposphere. In our model the widespread impact of bromine on tropospheric chemistry is caused by BrO mixing ratios that are around the detection limit of current instruments. For these reasons it is highly desirable to improve the detection limits, especially the sensitivity of space-borne instruments to BrO in the MBL to yield global coverage.

A first step to improve our knowledge about the global distribution of reactive bromine in the troposphere would be to conduct airborne studies in the free troposphere to check whether BrO is present at lower latitudes as well or if it is confined to high latitudes where it could already be directly measured.

An additional focus of future research in this field should be to investigate the contribution, spatial and temporal distribution of the different sources to global tropospheric bromine levels.

Acknowledgements. We thank C. Brühl for providing results from his 2D stratosphere model that we used to calculate the downward flux of bromine for the “strat” case. Many thanks to the reviewers, R. Sander, B.-M. Sinnhuber, and R. J. Salawitch for helpful comments on the ACPD version. This project was in part funded by the Deutsche Forschungsgemeinschaft DFG, Project GL353/1, MarHal.

Edited by: P. Monks

References

- Abbatt, J. P. D.: Heterogeneous reaction of HOBr with HBr and HCl on ice surfaces at 228 K, *Geophys. Res. Lett.*, 21, 665–668, 1994.
- Adams, J. W., Holmes, N. S., and Crowley, J. N.: Uptake and Reaction of HOBr on Frozen and dry Salt Surfaces, *Atmos. Chem. Phys.*, 2, 79–91, 2002, **SRef-ID: 1680-7324/acp/2002-2-79**.
- Aranda, A., Le Bras, G., La Verdet, G., and Poulet, G.: The BrO + CH_3O_2 reaction: Kinetics and role in the atmospheric ozone budget, *Geophys. Res. Lett.*, 24, 2745–2748, 1997.
- Atkinson, A., Baulch, D. L., Cox, R. A., Hampson, Jr., R. F., Kerr, J. A., Rossi, M. J., and Troe, J.: Evaluated kinetic and photochemical data for atmospheric chemistry: Supplement VI, *J. Phys. Chem. Ref. Data*, 26, 1329–1499, 1997.
- Atkinson, R., Baulch, D. L., Cox, R. A., Hampson, Jr., R. F., Kerr, J. A., Rossi, M. J., and Troe, J.: Summary of Evaluated Kinetic and Photochemical Data for Atmospheric Chemistry, Web Version, <http://www.iupac-kinetic.ch.cam.ac.uk>, 1999.
- Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Kerr, J. A., Rossi, M. J., and Troe, J.: Summary of Evaluated Kinetic and Photochemical Data for Atmospheric Chemistry, Web Version, Nov. 2003, <http://www.iupac-kinetic.ch.cam.ac.uk>, 2003.
- Bobrowski, N., Hönninger, G., Galle, B., and Platt, U.: Detection of bromine monoxide in a volcanic plume, *Nature*, 423, 273–276, 2003.
- Boucher, O., Moulin, C., Belviso, S., Aumont, O., Bopp, L., Cosme, E., von Kuhlmann, R., Lawrence, M. G., Pham, M., Reddy, M. S., Sciare, J., and Venkataraman, C.: Sensitivity study of dimethylsulphide (DMS) atmospheric concentrations and sulphate aerosol indirect radiative forcing to the DMS source representation and oxidation, *Atmos. Chem. Phys.*, 3, 49–65, 2003, **SRef-ID: 1680-7324/acp/2003-3-49**.

- Brasseur, G. P., Orlando, J. J., and Tyndall, G. S. (Eds.): *Atmospheric Chemistry and Global Change*, Oxford University Press, New York, Oxford, 1999.
- Brühl, C., Crutzen, P. J., and Groö, J.: High-latitude, summertime NO_x activation and seasonal ozone decline in the lower stratosphere: Model calculations based on observations by HALOE on UARS, *J. Geophys. Res.*, 103, 3587–3597, 1998.
- Crutzen, P. J. and Schmailzl, U.: Chemical budgets of the stratosphere, *Planet. Space. Sci.*, 31, 1009–1032, 1983.
- DeMore, W. B., Sander, S. P., Golden, D. M., Hampson, R. F., Kurylo, M. J., Howard, C. J., Ravishankara, A. R., Kolb, C. E., and Molina, M. J.: *Chemical Kinetics and Photochemical Data for Use in Stratospheric Modeling*, Tech. Rep. JPL Publication 97-4, Jet Propulsion Laboratory, Pasadena, CA, 1997.
- Dentener, F. and Crutzen, P. J.: Reaction of N_2O_5 on Tropospheric Aerosols: Impact on the Global Distributions of NO_x , O_3 , OH, *J. Geophys. Res.*, 98, 7149–7163, 1993.
- Fan, S.-M. and Jacob, D. J.: Surface ozone depletion in Arctic spring sustained by bromine reactions on aerosols, *Nature*, 359, 522–524, 1992.
- Fickert, S., Adams, J. W., and Crowley, J. N.: Activation of Br_2 and BrCl via uptake of HOBr onto aqueous salt solutions, *J. Geophys. Res.*, 104, 23 719–23 727, 1999.
- Fitzenberger, R., Bösch, H., Camy-Peyret, C., Chipperfield, M. P., Harder, H., Platt, U., Sinnhuber, B.-M., Wagner, T., and Pfeilsticker, K.: First Profile Measurements of Tropospheric BrO, *Geophys. Res. Lett.*, 27, 2921–2924, 2000.
- Harder, H., Camy-Peyret, C., Ferlemann, F., Fitzenberger, R., Hawat, T., Osterkamp, H., Schneider, M., Perner, D., Platt, U., Vradelis, P., and Pfeilsticker, K.: Stratospheric BrO profiles measured at different latitudes and seasons: Atmospheric observations, *Geophys. Res. Lett.*, 25, 3843–3846, 1998.
- Hausmann, M. and Platt, U.: Spectroscopic measurement of bromine oxide and ozone in the high Arctic during Polar Sunrise Experiment 1992, *J. Geophys. Res.*, 99, 25 399–25 413, 1994.
- Hebestreit, K., Stutz, J., Rosen, D., Matveiv, V., Peleg, M., Luria, M., and Platt, U.: DOAS Measurements of Tropospheric Bromine Oxide in Mid-Latitudes, *Science*, 283, 55–57, 1999.
- Hollwedel, J., Wenig, M., Beirle, S., Kraus, S., Köhl, S., Wilms-Grabe, W., Platt, U., and Wagner, T.: Year-to-Year Variability of Polar Tropospheric BrO as seen by GOME, *Adv. Space Res.*, 34, 804–808, 2004.
- Hönninger, G., Bobrowski, N., Palenque, E. R., Torrez, R., and Platt, U.: Reactive bromine and sulfur emissions at Salar de Uyuni, Bolivia, *Geophys. Res. Lett.*, 31, doi:10.1029/2003GL018 818, 2004.
- Hubinger, S. and Nee, J. B.: Absorption spectra of Cl_2 , Br_2 and BrCl between 190 and 600 nm, *J. Photochem. Photobiol. A: Chem.*, 86, 1–7, 1995.
- Ingham, T., Bauer, D., Sander, R., Crutzen, P. J., and Crowley, J. N.: Kinetics and Products of the Reactions $\text{BrO} + \text{DMS}$ and $\text{Br} + \text{DMS}$ at 298 K, *J. Phys. Chem. A*, 103, 7199–7209, 1999.
- Jefferson, A., Nicovich, J. M., and Wine, P. H.: Temperature-dependent kinetics studies of the reactions $\text{Br}(\text{}^2\text{P}_{3/2}) + \text{CH}_3\text{SCH}_3 \leftrightarrow \text{CH}_3\text{SCH}_2 + \text{HBr}$. Heat of formation of the CH_3SCH_2 radical, *J. Phys. Chem.*, 98, 7128–7135, 1994.
- Jungwirth, P. and Tobias, D. J.: Ions at the Air/Water Interface, *J. Phys. Chem. B*, 106, 6361–6373, 2002.
- Kettle, A. J., Andreae, M. O., Amouroux, D., Andreae, T. W., Bates, T. S., Berresheim, H., Bingemer, H., Boniforti, R., Curran, M. A. J., DiTullio, G. R., Helas, G., Jones, G. B., Keller, M. D., Kiene, R. P., Leck, C., Levasseur, M., Malin, G., Maspero, M., Matrai, P., McTaggart, A. R., Mihalopoulos, N., Nguyen, B. C., Novo, A., Putaud, J. P., Rapsomanikis, S., Roberts, G., Schebeske, G., Sharma, S., Simó, R., Staubes, R., Turner, S., and Uher, G.: A global database of sea surface dimethyl sulfide (DMS) measurements and a procedure to predict sea surface DMS as a function of latitude, longitude, and month, *Global Biogeochem. Cycles*, 13, 399–444, 1999.
- Langner, J. and Rodhe, H.: A global three-dimensional model of the tropospheric sulfur cycle, *J. Atmos. Chem.*, 13, 225–263, 1991.
- Lawrence, M. G., Crutzen, P. J., Rasch, P. J., Eaton, B. E., and Mahowald, N. M.: A model for studies of tropospheric photochemistry: Description, global distributions, and evaluation, *J. Geophys. Res.*, 104, 26 245–26 277, 1999.
- Lawrence, M. G., Jöckel, P., and von Kuhlmann, R.: What does the global mean OH concentration tell us?, *Atmos. Chem. Phys.*, 1, 37–49, 2001,
- SRef-ID: 1680-7324/acp/2001-1-37.**
- Leser, H., Hönninger, G., and Platt, U.: MAX-DOAS Measurements of BrO and NO_2 in the Marine Boundary Layer, *Geophys. Res. Lett.*, 30, 1537, doi:10.1029/2002GL015 811, 2003.
- Mallard, W. G., Westley, F., Herron, J. T., Hampson, R. F., and Frizzel, D. H.: *NIST Chemical Kinetics Database: Version 5.0*, National Institute of Standards and Technology, Gaithersburg, MD, 1993.
- McElroy, C. T., McLinden, C. A., and McConnell, J. C.: Evidence for bromine monoxide in the free troposphere during the Arctic polar sunrise, *Nature*, 397, 338–341, 1999.
- Orlando, J. J. and Tyndall, G. S.: Rate coefficients for the thermal decomposition of BrONO_2 and the heat of formation of BrONO_2 , *J. Phys. Chem.*, 100, 19 398–19 405, 1996.
- Platt, U. and Hönninger, G.: The role of halogen species in the troposphere, *Chemosphere*, 52, 325–338, 2003.
- Pundt, I., Van Roozendaal, M., Wagner, T., Richter, A., Chipperfield, M., Burrows, J. P., Fayt, C., Hendrick, F., Pfeilsticker, K., Platt, U., and Pommereau, J.-P.: Simultaneous UV-vis Measurements of BrO from Balloon, Satellite and Ground: Implications for Tropospheric BrO, in: *Proc. Fifth European Symp. on Polar Stratospheric Ozone 1999*, Air Poll. Res. Report 73, EUR 19340, European Commission, Brussels, Belgium, edited by: Harris, N. R. P., Guirlet, M., and Amanatidis, G. T., 316–319, 2000.
- Richter, A., Wittrock, F., Ladstätter-Weissenmayer, A., and Burrows, J. P.: GOME measurements of stratospheric and tropospheric BrO, *Adv. Space Res.*, 29, 1667–1672, 2002.
- Saiz-Lopez, A., Plane, J. M. C., and Shillito, J. A.: Bromine oxide in the mid-latitude marine boundary layer, *Geophys. Res. Lett.*, 31, doi:10.1029/2003GL018 956, 2004.
- Sander, R., Rudich, Y., von Glasow, R., and Crutzen, P. J.: The role of BrNO_3 in marine tropospheric chemistry: A model study, *Geophys. Res. Lett.*, 26, 2857–2860, 1999.
- Sander, S. P., Friedl, R. R., Golden, D. M., Kurylo, M. J., Huie, R. E., Orkin, V. L., Moortgat, G. K., Ravishankara, A. R., Kolb, C. E., Molina, M. J., and Finlayson-Pitts, B. J.: *Chemical Kinetics and Photochemical Data for Use in Stratospheric Modeling*, Tech. Rep. JPL Publication 02-25, Jet Propulsion Laboratory, Pasadena, CA, 2003.

- Singh, H. B. and Zimmerman, P. B.: Atmospheric distribution and sources of nonmethane hydrocarbons, *Adv. Environ. Sci. Technol.*, 24, 177–235, 1992.
- Stutz, J., Hebestreit, K., Alicke, B., and Platt, U.: Chemistry of Halogen Oxides in the Troposphere: Comparison of Model Calculations with Recent Field Data, *J. Atmos. Chem.*, 34, 65–85, 1999.
- Stutz, J., Ackermann, R., Fast, J. D., and Barrie, L.: Atmospheric Reactive Chlorine and Bromine at the Great Salt Lake, Utah, *Geophys. Res. Lett.*, 29, doi:10.1029/2002GL014812, 2002.
- Toumi, R.: BrO as a sink for dimethylsulphide in the marine atmosphere, *Geophys. Res. Lett.*, 21, 117–120, 1994.
- Van Roozendaal, M., Wagner, T., Richter, A., Pundt, I., Arlander, D. W., Burrows, J. P., Chipperfield, M., Fayt, C., Johnston, P. V., Lambert, J.-C., Kreher, K., Pfeilsticker, K., Platt, U., Pommereau, J.-P., Sinnhuber, B.-M., Tørnkvist, K. K., and Wittrock, F.: Intercomparison of BrO measurements from ERS-2 GOME, ground-based and balloon platforms, *Adv. Space Res.*, 29, 1661–1666, 2002.
- Vogt, R., Crutzen, P. J., and Sander, R.: A mechanism for halogen release from sea-salt aerosol in the remote marine boundary layer, *Nature*, 383, 327–330, 1996.
- von Glasow, R. and Crutzen, P. J.: Tropospheric halogen chemistry, in: *The Atmosphere*, edited by: Keeling, R. F., Vol. 4, *Treatise on Geochemistry*, edited by: Holland, H. D. and Turekian, K. K., pp. 21–64, Elsevier-Pergamon, Oxford, 2003.
- von Glasow, R. and Crutzen, P. J.: Model study of multiphase DMS oxidation with a focus on halogens, *Atmos. Chem. Phys.*, 4, 589–608, 2004, **SRef-ID: 1680-7324/acp/2004-4-589**
- von Glasow, R., Sander, R., Bott, A., and Crutzen, P. J.: Modeling halogen chemistry in the marine boundary layer. 1. Cloud-free MBL, *J. Geophys. Res.*, 107, 4341, doi:10.1029/2001JD000942, 2002a.
- von Glasow, R., Sander, R., Bott, A., and Crutzen, P. J.: Modeling halogen chemistry in the marine boundary layer. 2. Interactions with sulfur and cloud-covered MBL, *J. Geophys. Res.*, 107, 4323, doi:10.1029/2001JD000943, 2002b.
- von Kuhlmann, R., Lawrence, M. G., Crutzen, P. J., and Rasch, P. J.: A Model for Studies of Tropospheric Ozone and Non-Methane Hydrocarbons: Model Description and Ozone Results, *J. Geophys. Res.*, 108, 4294, doi:10.1029/2002JD002893, 2003.
- Wagner, T. and Platt, U.: Satellite mapping of enhanced BrO concentrations in the troposphere, *Nature*, 395, 486–490, 1998.
- Wagner, T., Leue, C., Wenig, M., Pfeilsticker, K., and Platt, U.: Spatial and temporal distribution of enhanced boundary layer BrO concentrations measured by the GOME instrument aboard ERS-2, *J. Geophys. Res.*, 106, 24 225–24 235, 2001.
- WMO: Scientific assessment of ozone depletion: 2002, World Meteorological Organization, Global Ozone Research and Monitoring project, Report Nr. 47, 2003.